

A central graphic features a red human heart with white veins, set against a dark blue background with white stars. The heart is surrounded by a cluster of stylized, colorful geometric shapes resembling shards of broken glass or shards of ice, in shades of red, white, and blue. The overall aesthetic is modern and abstract.

Schizotypal Diorder: Management Challenges with **Clozapine-induced Myocarditis**

Ahmed Karim, MD


PGY3 Psychiatry Resident

Disclosure

I currently have no relationships of any kind with any company whose products or services in any way relate to the practice of medicine, medical education or research.

Goals of Presentation

- Introduction to Personality Disorders
- Differentiate between Schizophrenia and Schizotypal Personality Disorder SPD.
- Differentiate Autism from Schizotypal Personality Disorder SPD.
- Roles of Antipsychotics in Schizotypal Personality Disorder.
- Review the Diagnostic Criteria and Treatment Options.
- Exploring Diagnostic Criteria and Etiologies for Clozapine-Induced Myocarditis (CIM).




Personality disorders

- **Definition:** Inflexible, maladaptive, and pervasive behavior patterns causing distress or impaired functioning.
 - Typically egosyntonic (feels okay to me!) and present by early adulthood.

Clusters:

- **Cluster A ("Weird"):** Odd or eccentric behaviors; difficulty forming social relationships. Linked with schizophrenia.
 - Paranoid: Pervasive distrust, suspicion, and hypervigilance. Individuals often have a cynical view of the world.
 - Schizoid: socially withdrawn and solitary activities. Displays limited emotional expression and indifference to others' opinions.
 - Schizotypal: Marked by eccentric appearance, odd beliefs, and interpersonal awkwardness. Associated with the schizophrenia spectrum.



(Le et al., 2024, p. 582)

Personality disorders

- **Cluster B ("Wild"):** Dramatic, emotional, or erratic behaviors.
 - Associated with mood disorders and substance use.
 - Antisocial: Disregard for others, criminality, impulsivity, no remorse; linked to conduct disorder before age 15; more common in males.
 - Borderline: Unstable mood, relationships, self-image; fear of abandonment, impulsivity, self-harm; dialectical behavior therapy often used.
 - Histrionic: Attention-seeking, dramatic, shallow emotions, provocative behavior; often uses appearance to gain attention.
 - Narcissistic: Grandiosity, need for admiration, lack of empathy; reacts defensively to criticism, envious of others.



Personality disorders


- **Cluster C ("Worried"):** Anxious or fearful behaviors. Associated with anxiety disorders
 - Avoidant: Very sensitive to rejection and criticism, avoids social situations, but still wants relationships despite feeling inadequate.
 - Obsessive-Compulsive: Focuses on order, perfection, and control; actions match their beliefs.
 - Dependent: Needs constant support and reassurance, lacks self-confidence, may stay in harmful relationships.



Feature	Schizotypal Personality Disorder (STPD)	Schizophrenia/Psychosis
Main Traits	Odd beliefs, strange thinking, and social anxiety	Delusions, hallucinations, and jumbled thinking
When It Starts	Teenage years or early adulthood	Late teenage years to early adulthood
How Long It Lasts	Long-term, relatively steady symptoms	Long-term, with sudden intense episodes of psychosis
Thinking Ability	Mildly affected; thoughts may be odd or unusual	Often seriously affected; thinking can be confused or slow
Social Life	Ongoing “ intense ” social anxiety, not many close friends	Major problems in social and work life
Medication Response	Medication may NOT be very effective; usually not needed unless psychotic episodes happen	Antipsychotic medications are the main treatment
Paranoia	Mild to moderate; often linked to social anxiety	Can be severe; paranoid thoughts are common

Patient Demographics/Identifying Data:

- **Age:** Late twenties
- **Gender:** Male
- **Psychiatric History:** Schizotypal Personality Disorder
- **Legal:** 1st degree relative is the guardian. He was in detention center.
Referred for forensic evaluation. Opined CST and released on Oct 2022.




Substance Use History

- **Past Use:** Cannabis, LSD, mushrooms
- **Last Use:** 2017 (episodes started in 2017 after heavy use of LSD and mushrooms.)
- **Connection to Symptoms:** Patient attributes telepathic voices to LSD


Social History

- **Living Situation** With mother (guardian) History of witnessing domestic violence
- **Relationships** None due to mental illness
- **Employment** Unemployed
- **Education** High school graduate



Past Psychiatric History

- **Multiple Admissions:** August 2022, April 2023, August 2023, March 2024 and April 2024
- **Forensic Admission:** Competency restoration, diagnosed with Schizotypal PD, unspecified psychosis, cannabis use disorder
- **Treatment History:** Various antipsychotics (limited effect), stabilized on Diazepam and Sertraline, DBT participation
- Multiple admissions for self-harm and aggression
- **Diagnosis** of schizotypal personality disorder, unspecified psychosis, cannabis use disorder
- History of impulsivity, anger, and self-injury
- **Medications Trials:** of paliperidone, Depakote, aripiprazole, and naltrexone with limited response
- Improvement on diazepam and sertraline



Forensic psychiatric evaluation for competency to stand trial in Jan 2022

• **Primary Discharge Diagnosis:** Schizotypal Personality Disorder.

- **Secondary Diagnoses:** Unspecified Psychosis & Cannabis Use Disorder.

• **Symptomology:**

- Odd beliefs or magical thinking influencing behavior.
- Overelaborate, vague, or metaphorical speech.
- Odd or eccentric behaviors.
- Social isolation.
- Odd affect.

• **Behavioral Patterns During Hospitalization:**

- Impulsivity, anger, and self-injury (e.g., hitting himself in the head).
- Emotional dysregulation, including screaming and verbal aggression towards staff.
- Frequent use of lorazepam 2 mg PRN for emotional outbursts.

•Medication Trials:

- Paliperidone, Valproate (up to 1,500 mg/day), Aripiprazole, and Naltrexone—NO meaningful response.

•Effective Medication Regimen:

- **Diazepam 5 mg BID:** Reduced self-harming behaviors and emotional outbursts.
- **Sertraline (up to 200 mg/day):** Improved isolation, anxiety, and depression.
- **Melatonin 6 mg QHS:** Managed insomnia.
- **Therapy:** Participated in Dialectical Behavior Therapy (DBT).
- **Competency Status:** Opined competent to stand trial August 2022. Discharged to the County Detention Center.

•Discharge Medications:

- Diazepam 5 mg BID (scheduled) for impulsivity, severe anxiety, and self-harm.
- Sertraline 200 mg for anxiety and depression.
- Melatonin 6 mg QHS for insomnia.

(Oct 2022)

- **Initial Presentation/Reason for admission:**

- Nonadherent to medications
- After 3 days, he was discharged on parole to the care of his mother. But, he was not given a 30-day supply
- Self-injurious behavior post-release from detention. He felt others were provoking him, leading to frustration and scratching his face.
- Involuntary admission (96-hour hold)

- **Key Symptoms:**

- Self-harm (scratching face, hitting head), fixated on vaccine indications, intervals, and pathophysiology.

- **Mental Status Exam:** Coherent, cooperative


- Denies SI/HI/AVH
- Talked to himself and exhibited odd behaviors (e.g., screaming at the mirror, yelling in his room).

- **Legal:** relative (1st Degree) was seeking guardianship


- **Length of stay:** 41 days

- **Outcome:**

- Consistently denied SI, HI and AVH.
- No aggressive or self-injurious behaviors observed.
- Restarted on Valium 5 mg BID, Zoloft 200 mg daily, and slept well with trazodone.
- QMHP worked on level 2 placement; mother decided to bring him home.



Feature	Odd Beliefs	Delusions
What It Is	Unusual beliefs that seem strange but could be possible.	Strongly held false beliefs that aren't true, even with proof.
Connection to Reality	Usually aware that others don't share the belief.	Firmly believe it's true, even when proven wrong.
Examples	Believing in lucky charms or strange superstitions.	Thinking the government is spying on you without any evidence.
Cultural Influence	Influenced by culture, religion, or community.	May start from cultural ideas but become extreme and irrational.
Flexibility	Can change or be dropped if proven wrong.	Hard to change; belief stays strong even with evidence against it.



(Summer 2023)

- **Reason for admission**

- admitted after assaulting his mother and peers
- Pt requested to be taken to hospital
- Prior to that he spent 2-3 days in jail following incidents
- His body being controlled from outside by multiple people

- **Psychological Assessment**

- Schizotypal Personality Disorder based on testing: Minnesota Multiphasic Personality Inventory

- **Incident**

- Aggressive episode – he requested lorazepam, became agitated, attempted to kick and spit on staff. Manual hold applied

- **Length of stay:** 101 days

- **Medication Adjustment**


- Transitioned from olanzapine to loxapine 10 mg twice daily (increased to 10 mg AM and 30 mg PM)

Treatment Plan:

- (From Outpatient) Olanzapine 10 mg twice daily for psychosis and aggression.
- Sertraline 200 mg daily for mood management (anxiety). Increase loxapine from 10mg to 20mg BID for psychosis
- Gradually tapered off from Olanzapine and started loxapine 10 mg BID that was gradually increased to 10 mg in QAM and 30 mg at QHS.
- Reported not hearing voices, a telepathic communications are better pleasant in nature.


On Discharge:

- He was observed smiling to himself
- Discharge to the mother



Diagnostic Considerations

- **Autism Spectrum Disorder (consider social deficits)**
 - Is there evidence suggesting that autistic traits are associated with schizotypal traits?
 - A chinses study examined the factor structure of the Autism Spectrum Quotient 10 (AQ-10) and its associations with schizotypal traits (measured by the Schizotypal Personality Questionnaire-Brief [SPQ-B])
 - There is evidence suggesting that autistic traits are associated with schizotypal traits. This study examined the factor structure of the Autism Spectrum Quotient 10 (AQ-10) and its associations with schizotypal traits (measured by the Schizotypal Personality Questionnaire-Brief [SPQ-B])



AQ-10

Autism Spectrum Quotient (AQ)

A quick referral guide for adults with suspected autism who do not have a learning disability.

Please tick one option per question only:


	Definitely Agree	Slightly Agree	Slightly Disagree	Definitely Disagree
--	------------------	----------------	-------------------	---------------------

1 I often notice small sounds when others do not				
2 I usually concentrate more on the whole picture, rather than the small details				
3 I find it easy to do more than one thing at once				
4 If there is an interruption, I can switch back to what I was doing very quickly				
5 I find it easy to 'read between the lines' when someone is talking to me				
6 I know how to tell if someone listening to me is getting bored				
7 When I'm reading a story I find it difficult to work out the characters' intentions				
8 I like to collect information about categories of things (e.g. types of car, types of bird, types of train, types of plant etc)				
9 I find it easy to work out what someone is thinking or feeling just by looking at their face				
10 I find it difficult to work out people's intentions				

SCORING: Only 1 point can be scored for each question. Score 1 point for Definitely or Slightly Agree on each of items 1, 7, 8, and 10. Score 1 point for Definitely or Slightly Disagree on each of items 2, 3, 4, 5, 6, and 9. If the individual scores **more than 6 out of 10**, consider referring them for a specialist diagnostic assessment.

This test is recommended in 'Autism: recognition, referral, diagnosis and management of adults on the autism spectrum' (NICE clinical guideline CG142). www.nice.org.uk/CG142

Key reference: Allison C, Auyeung B, and Baron-Cohen S. (2012) *Journal of the American Academy of Child and Adolescent Psychiatry* 51(2):202-12.




Autistic traits were positively correlated with schizotypal traits of disorganized features ($r = 0.21$, $p < 0.001$), interpersonal relationship deficits ($r = 0.19$, $p < 0.001$), and cognitive-perceptual deficits ($r = 0.11$, $p = 0.001$).




• Autism Spectrum Disorder: DSM-V.

- 1. Social Deficits in Conversation and Relationships:**
- 2. Sensory Sensitivities:**
- 3. Difficulty Making Eye Contact:**
- 4. Dislike of Change:**
- 5. Hyperfixation on Certain Interests:**
- 6. Repetitive Behaviors:**




- Stanfield et al., 2017 – was This is the first fMRI study directly comparing Autism Spectrum Disorder (ASD) and Schizotypal Personality Disorder (SPD).
- Clinical symptoms were almost similar.
- SPD shows hyperactivation in social brain regions (amygdala, cerebellum, fusiform gyrus) compared to ASD.
- Long story short, ASD =/≠ SPD.




(March 2024)

- Self-harm, aggression towards mother and dogs
- Similar to previous presentations with linear thought process and calm demeanor
- Continues to discuss telepathic communications and mind control
- Exacerbated anxiety? moving, mother's job loss, and leaving a friend


- **Diagnoses**
 - Schizotypal personality disorder
 - Unspecified psychotic disorder
 - History of cannabis use disorder
- **Treatment plan**
 - Start Clozapine titratic




Treatment Course and Test Results



- 1st week since admission: Clozapine started on 1st of March and titrated up to 50mg AM and 100mg PM
- 2nd week since admission: He experienced low grade fever with T 100.1°F, with headache, nausea, and myalgias. He reports that this has completely resolved.
 - CBC, CRP, troponin and EKG (QTC 283/423)
 - Slightly elevated troponin of 5.3.
 - Denied chest pain, shortness of breath, irregular heartbeat, dizziness, or headache.
 - Received a COVID vaccine on 03/13/24.
- Troponin levels at **15.8** (greater than twice the upper limit of normal-Normal high 4.9) , which was trending upward from **5.3** to **14.1**, alongside normal CRP levels (12.1).
- In the ED:
 - -Negative Sed rate, downtrending CRP (9.8), and troponin (6.5).




- 3rd week since admission:
 - 2nd ED admission
 - Troponin levels in 2 days:
 - **48.5 pg/ml**
 - **67.9 pg/ml**
 - **110.5 pg/ml**
 - **162.7 pg/ml**
 - **664.1 pg/ml**
 - **562.8 pg/ml**
 - CRP 2 days:
 - **38.9**
 - **45.7**
 - **70.7**




Cardiac Workup

- Serial of EKG
- US Echocardiogram –
 - Left ventricular systolic function is mildly reduced, estimate is 45-50%.
 - View suggested distal anterior and apical hypokinesis
 - Diastolic function could not be accurately assessed due to tachycardia.
 - The right ventricle is normal
- 3 days after the echo, Cardiac Nuclear Stress Test performed:
 - Myocardial perfusion imaging is normal. There is no evidence of ischemia.
 - LVEF is 64%. Post stress wall motion is normal.
- Patient hemodynamically stable and asymptomatic
- Negative sed rate, downtrending CRP and troponin
- No indication for further workup or monitoring at that time
- Transthoracic echocardiogram showed mildly reduced LV systolic function and possible distal anterior and apical hypokinesis
- Plan to schedule for outpatient cardiac MRI to further investigate possible myocarditis




Vent. Rate Over Time



Vent Rate: 60 - 100 bpm

normal range of the QT/QTc: 350 - 470 ms


QTc Over Time



Signs and symptoms in literature:

- 87% of patients develop symptoms within the first month of treatment.
- The median age of patients and dose of clozapine at presentation was 30 years and 250mg/day respectively
- Clinical presentation included:
 - Shortness of breath (67%)
 - Fever (67%)
 - Tachycardia (58%).
 - Cardiac markers are elevated in 87% of the 54 cases that reported these markers.
 - Global ventricular dysfunction was the predominant echocardiogram finding (57%).

(Bellissima, Tingle et al. 2018)



(Adapted from: Jeong, S. H., Ahn, Y. M., Koo, Y. J., et al. (2002). The characteristics of clozapine-induced fever. *Schizophrenia Research*, 56, 191-193)

Presenting features of adverse reactions to clozapine and of biopsy-proven idiopathic myocarditis


Clozapine adverse drug reactions cited by researchers, in order of frequency⁶	Presenting features in patients with biopsy-proven idiopathic myocarditis¹³
Electro- or echocardiographic abnormality (66%)	Abnormal ECG in all patients (prolonged QT interval in 90%)
Fever (49%)	Dyspnoea (90%)
Tachycardia (46%)	Palpitations/arrhythmia (70%)
Elevated troponin level (36%)	Elevated levels of cardiac biomarkers (70%)
Chest pain (32%)	Influenza-type illness/viral prodrome (including fever) (50%)
Elevated creatine kinase level (31%)	Tachycardia (40%)
Leucocytosis (28%)	Leucocytosis (35%)
Dyspnoea (27%)	Chest pain (25%)

ECG = electrocardiogram.

(Ramamurthy et al., 1993)

Etiology:


- The mechanisms remain unclear.
- leading hypothesis is that of an IgE-mediated hypersensitivity reaction.
 - Supported peripheral eosinophilia and eosinophilic inclusions within endomyocardial biopsy sample.
- Clozapine had higher noradrenaline which may lead to left ventricular (LV) dysfunction
 - Evidence suggests increase catecholamines may cause cardiac dysfunction. (Elman et al., 1999)
 - For example, increased plasma catecholamine levels have been implicated in Takotsubo cardiomyopathy (reversible form of LV dysfunction. Thus it is plausible that increased catecholamine levels could be a contributing factor in the development of myocarditis and/or cardiomyopathy among affected patients. (Novak et al., 2007)
- The odds of clozapine-induced myocarditis increased with concurrent sodium valproate use (Ronaldson et al., 2012)
 - (k = 6, n = 903, pooled OR 3.58, 95% CI 1.81–7.06)
- Other unproven mechanisms (Devarajan et al., 2000)
 - Cytochrome P450 1A2/1A3 enzyme deficiencies
 - Blockade of calcium-dependent ion channels
 - Increased production of inflammatory cytokines
 - Low serum selenium levels.



(Elman et al., 1999)


Low Selenium

- According to some studies, clozapine (Clozaril) can lower blood levels of selenium
 - Selenium concentrations in plasma and red cells → significantly lower in schizophrenic patients treated with clozapine as compared with all other groups. (Vaddadi, E. Soosai, and G. Vaddadi 2003)
 - “The reactive metabolite (nitrenium)of clozapine reacts with glutathione, and therefore, it is likely that it also binds to selenocysteine-containing proteins, such as glutathione peroxidase, thioredoxin reductase, and protein disulfide isomerase.” (Ip & Utrecht, 2008)
 - The highest in myelocyte stage (neutrophil myelocytes?)
 - Clozapine is activated to free radical metabolites; activated neutrophils may metabolize clozapine to a reactive nitrenium ion, thereby producing agranulocytosis by either direct toxicity or an immune-mediated mechanism. (Linday et al., 1995)




Brain Imaging:

- April 2022 CT Head w/o
 - Cont No acute intracranial abnormality.
- April, 2024 Head w/o Cont
 - Right cheek soft tissue swelling/hematoma.



- **Question:** Which brain structure is primarily involved in the consolidation of declarative memory (facts and events)?
 - A) Prefrontal cortex and amygdala
 - B) Parahippocampal gyrus
 - C) Basal ganglia
 - D) Thalamus and hypothalamus




Neuro-Imaging findings in studies?

- Brain images (**CT/MRI**) in SPD show abnormalities in the superior temporal gyrus, parahippocampus, and corpus callosum, like schizophrenia, but without medial temporal lobe or lateral ventricle abnormalities. (Dickey et al., 2002)
 - With medial temporal lobe.? (Dickey et al., 1999)

fMRI


- Individuals with SPD show functional alterations in the **striatal, frontal, and temporal brain areas**, which align with findings observed across the schizophrenia spectrum. (Chrysoula Zouraraki et al., 2023)
- **Schizotypy and Social Cognition:** Negative schizotypy (e.g., anhedonia) is linked to reduced brain activity in regions associated with theory of mind (ToM) and empathy (Wang et al., 2015).



(Wang et al., 2015).

Anti-psychotic medication role in SPD?

- Minimal evidence regarding pharmacotherapy (Koch et al., (2016) 6 (2): 75–81.)
 - Olanzapine, risperidone, haloperidol, fluoxetine, and thiothixene showed benefits, but such agents should be considered on a case-by-case basis. (Koch et al., (2016) 6 (2): 75–81.)
- Risperidone was the most often studied drug class and were described as beneficial. Studies on the longitudinal course described a moderate remission rate and possible conversion rates to other schizophrenia spectrum disorders. (Kirchner et al., 2018)
- Risperidone appears to be effective in reducing symptom severity in SPD and is generally well tolerated. (Koenigsberg et al., 2003)



(April 2024)

- Self-harm, aggression towards mother and dogs
- Similar to previous presentations with linear thought process and calm demeanor
- Continues to discuss telepathic communications and mind control
- Open about stressors: moving, mother's job loss, and leaving a friend
- History of witnessing domestic violence and experiencing a severe dog attack

Medications course:

- Loxapine, titrated up to 100 mg PO daily.
- Added Cariprazine 1.5 mg PO QD as dual therapy, then increased to 6 mg PO daily.

Adverse Effects and Adjustments:

- Loxapine was discontinued due to (EPS).
- Cariprazine monotherapy was discontinued at 6 mg PO QD after 3 weeks, due to a lack of symptom improvement.

New Treatment Plan:

- Patient reported previous improvement on Risperidone monotherapy.
- Risperidone was started and titrated to 3 mg PO daily, which is the discharge dose.



Additional Medications:

- While on Cariprazine monotherapy, the patient requested PRN Lorazepam (generic name for Ativan) 1-2 times per day, reporting that it helped quiet telepathic communication.
- After switching to Risperidone, the patient reduced PRN Lorazepam usage and has not requested it for 5 days prior to discharge.

Treatment for PTSD Symptoms:

- Sertraline 50 mg PO QD then increased to 150 mg PO daily.

Length of Stay Chart



Mortality sequelae:


- The mortality rate associated with clozapine-induced myocarditis varies across studies, with reported rates generally ranging from around 10% to as high as 50%.
 - **24% and 64%** (Kilian et al., 1999; Remington et al., 2017). In a **6** year long prospective cohort study with 503 patients on clozapine in Australia, 3% developed myocarditis and 2% had sudden death with 90% of the deaths attributed to cardiac etiologies including myocarditis, cardiomyopathy, and idiopathic arrhythmias (Khan et al., 2017).
 - **10% to 46%** (Sackey et al. 2018)
 - **10%** - Ronaldson, K. J., et al. (2010). "Rapid clozapine titration and increased risk of clozapine-induced myocarditis." *The Journal of Clinical Psychiatry*, 71(11), 1566-1571 .
 - **10% to 20%** - Bellissima, B. L., et al. (2018). "The incidence of clozapine-induced myocarditis: A systematic review and meta-analysis." *Journal of Clinical Psychopharmacology*, 38(5), 451-458 .
 - Around **20%** - Killick, S. K., et al. (2018). "Clozapine-induced myocarditis: A case series and literature review." *General Hospital Psychiatry*, 54, 45-49 .
 - **12-13%** - Reinders, J., et al. (2020). "Clozapine-induced myocarditis: An analysis of 38 cases." *Australian & New Zealand Journal of Psychiatry*, 44(1), 63-67 .

(Li, Gurrera, and Delisi 2018)

Feature	Schizotypal Personality Disorder (STPD)	Schizophrenia/Psychosis
Main Traits	Odd beliefs, strange thinking, and social anxiety	Delusions, hallucinations, and jumbled thinking
When It Starts	Teenage years or early adulthood	Late teenage years to early adulthood
How Long It Lasts	Long-term, relatively steady symptoms	Long-term, with sudden intense episodes of psychosis
Thinking Ability	Mildly affected; thoughts may be odd or unusual	Often seriously affected; thinking can be confused or slow
Social Life	Ongoing “ intense ” social anxiety, not many close friends	Major problems in social and work life
Medication Response	Medication may NOT be very effective; usually not needed unless psychotic episodes happen	Antipsychotic medications are the main treatment
Paranoia	Mild to moderate; often linked to social anxiety	Can be severe; paranoid thoughts are common

Take home points

1. Diagnosing SPD can be challenging and requires a multidisciplinary team approach.
2. Consider managing anxiety as a key component of treatment.
3. Monitor vitals closely, especially during the first weeks.
4. Troponin and CK levels are sensitive indicators for detecting CIM.
5. Therapy and behavioral interventions are crucial alongside pharmacotherapy.



Reference list:

- Le, T., Bhushan, V., Qiu, C., Chalise, A., & Kaparaliotis, P. (2024). *First Aid for the USMLE Step 1 2024: A Student-to-Student Guide*. McGraw-Hill Education.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing. Available at <https://www.psychiatry.org/psychiatrists/practice/dsm>
- National Institute of Mental Health. (n.d.). Schizophrenia. Retrieved from <https://www.nimh.nih.gov/health/topics/schizophrenia/index.shtml>
- Mayo Clinic. (n.d.). Schizophrenia. Retrieved from <https://www.mayoclinic.org/diseases-conditions/schizophrenia/symptoms-causes/syc-20354443>
- Mayo Clinic. (n.d.). Schizotypal Personality Disorder. Retrieved from <https://www.mayoclinic.org/diseases-conditions/schizotypal-personality-disorder/symptoms-causes/syc-20353919>
- MedlinePlus. (n.d.). Schizophrenia. U.S. National Library of Medicine. Retrieved from <https://medlineplus.gov/schizophrenia.html>
- MedlinePlus. (n.d.). Personality Disorders. U.S. National Library of Medicine. Retrieved from <https://medlineplus.gov/personalitydisorders.html>
- Bess Yin-Hung, L., et al. (2024). "Factor analysis of the Chinese version of the Autism Spectrum Quotient 10 and its association with schizotypal traits in adolescents and young adults in Hong Kong." *East Asian Archives of Psychiatry* 34(2): 23-28.
- Vickers, M., et al. (2022). "Risk factors for clozapine-induced myocarditis and cardiomyopathy: A systematic review and meta-analysis." *Acta psychiatrica Scandinavica* 145(5): 442-455.
- Bellissima, B. L., et al. (2018). "A systematic review of clozapine-induced myocarditis." *Int J Cardiol* 259: 122-129.
- Novak, G., Kross, K., Follmer, K., Brofferio, A., & Jamshid Shirani. (2007). Transient Biventricular Apical Ballooning: A Unique Presentation of the "Broken Heart." *Clinical Cardiology*, 30(7), 355–358. <https://doi.org/10.1002/clc.20065>
- Devarajan, S., Kutcher, S., & SM Dursun. (2000). Clozapine and sudden death. *The Lancet*, 355(9206), 841–841. [https://doi.org/10.1016/s0140-6736\(00\)00035-0](https://doi.org/10.1016/s0140-6736(00)00035-0)
- Elman, I., Goldstein, D. S., Graeme Eisenhofer, Folio, J., Malhotra, A. K., Adler, C. M., Pickar, D., & Breier, A. (1999). Mechanism of Peripheral Noradrenergic Stimulation by Clozapine. *Neuropsychopharmacology*, 20(1), 29–34. [https://doi.org/10.1016/s0893-133x\(98\)00047-5](https://doi.org/10.1016/s0893-133x(98)00047-5)
- Ramamurthy, S., Talwar, K. K., Goswami, K. C., Shrivastava, S., Chopra, P., S Broor, & Malhotra, A. (1993). Clinical profile of biopsy proven idiopathic myocarditis. *International Journal of Cardiology*, 41(3), 225–232. [https://doi.org/10.1016/0167-5273\(93\)90119-2](https://doi.org/10.1016/0167-5273(93)90119-2)

Reference list:

- Li, Kevin J, Ronald J Gurrera, and Lynn E Delisi. 2018. "Potentially Fatal Outcomes Associated with Clozapine." *Schizophrenia Research* 199 (September): 386–89. <https://doi.org/10.1016/j.schres.2018.02.058>.
- Sackey, Bryan K, Troy A Moore, Nicole L Cupples, and Cynthia A Gutierrez. 2018. "Clozapine-Induced Myocarditis: Two Case Reports and Review of Clinical Presentation and Recognition." *Mental Health Clinician* 8 (6): 303–8. <https://doi.org/10.9740/mhc.2018.11.303>.
- Vaddadi, K S, E. Soosai, and G. Vaddadi. 2003. "Low Blood Selenium Concentrations in Schizophrenic Patients on Clozapine." *British Journal of Clinical Pharmacology* 55 (3): 307–9. <https://doi.org/10.1046/j.1365-2125.2003.01773.x>.
- Ip, J., & Utrecht, J. P. (2008). Testing the hypothesis that selenium deficiency is a risk factor for clozapine-induced agranulocytosis in rats. *Chemical Research in Toxicology*, 21(4), 896-905. <https://doi.org/10.1021/tx7004037>
- Lindsay, L. , Pippenger, , Howard, A. & Lieberman, J. (1995). Free Radical Scavenging Enzyme Activity and Related Trace Metals in Clozapine-Induced Agranulocytosis. *Journal of Clinical Psychopharmacology*, 15 (5), 353-360.
- Lindsay, L. A., Pippenger, C. E., Howard, A. M., & Lieberman, J. A. (1995). Free radical scavenging enzyme activity and related trace metals in clozapine-induced agranulocytosis: A pilot study.
- Wang, Y., et al. (2015). Dimensional schizotypy and social cognition: an fMRI imaging study. *Front. Behav. Neurosci.*
- Cannon, T. D., et al. (2009). Structural abnormalities in schizotypal personality disorder. *Psychiatry Research: Neuroimaging*.
- Van der Meer, L., et al. (2010). Emotion regulation in schizophrenia. *Schizophrenia Bulletin*.
- Lu, C.-P., et al. (2023). Cognitive and affective empathy in schizophrenia. *Schizophrenia Research*.
- Koenigsberg, H. W., Reynolds, D., Goodman, M., New, A. S., Mitropoulou, V., Trestman, R. L., & Siever, L. J. (2003). Risperidone in the Treatment of Schizotypal Personality Disorder. *The Journal of Clinical Psychiatry*, 64(6), 18624. <https://www.psychiatrist.com/jcp/risperidone-treatment-schizotypal-personality-disorder/>
- Stanfield, A. C., Philip, R. C. M., Whalley, H., Romaniuk, L., Hall, J., Johnstone, E. C., & Lawrie, S. M. (2017). Dissociation of brain activation in autism and schizotypal personality disorder during social judgments. *Schizophrenia Bulletin*, 43(6), 1220-1230. <https://doi.org/10.1093/schbul/sbx097>
- Chrysoula Zouraraki, Karamaouna, P., & Giakoumaki, S. G. (2023). Cognitive Processes and Resting-State Functional Neuroimaging Findings in High Schizotypal Individuals and Schizotypal Personality Disorder Patients: A Systematic Review. *Brain Sciences*, 13(4), 615–615. <https://doi.org/10.3390/brainsci13040615>
- Dickey, C. C., McCarley, R. W., & Shenton, M. E. (2002). The Brain in Schizotypal Personality Disorder: A Review of Structural MRI and CT Findings. *Harvard Review of Psychiatry*, 10(1), 1–15. <https://doi.org/10.1080/10673220216201>
- Dickey, C. C., McCarley, R. W., Voglmaier, M. M., Niznikiewicz, M. A., Seidman, L. J., Yoshio Hirayasu, Fischer, I., Eng Kaet Teh, Richard Van Rhoads, Jakab, M., Kikinis, R., Jolesz, F. A., & Shenton, M. E. (1999). Schizotypal personality disorder and MRI abnormalities of temporal lobe gray matter. *Biological Psychiatry*, 45(11), 1393–1402. [https://doi.org/10.1016/s0006-3223\(99\)00030-x](https://doi.org/10.1016/s0006-3223(99)00030-x)